

# 2022 Pediatric Open Fracture Evaluation Clinical Guideline

PED

**Start**

Overlying break in the skin or concern for open fracture?

YES

Hemodynamically Stable

YES

**Stabilize and Observe**

YES

- Neurovascular assessment
- Check tetanus immunization status and order vaccine and/or immunoglobulin as needed

**Obtain images**

Is bedside orthopedic consult available within 60 min?

YES

Orthopedic Surgery Consult

NO

**Transfer to PED+ or PED1**

**Start antibiotics prior to transfer\*\***

NO

**Transfer to PED1**

While awaiting transfer, important to assess and initiate resuscitation

- Fluid resuscitation
- Packed RBCs if indicated
- Assess for other causes of bleeding
- Start antibiotics prior to transfer\*\*

**Don't delay transport to obtain imaging**

**Time Goal: Identify open fractures within 30 min of arrival**

**Time Goal: Start antibiotics within 60 min of arrival**

This algorithm does not replace clinical judgment and is not intended to be prescriptive for all patients

**\*\*See table 1 and 2 for classification and recommendation)**

# 2022 Pediatric Open Fracture Evaluation Clinical Guideline

**Table 1: Gustilo Classification**

<b>Gustilo Grade</b>	<b>Definition</b>
I	Open fracture, clean wound, wound < 1 cm in length
II	Open fracture, wound > 1 cm but < 10 cm in length without extensive soft-tissue damage, flaps or avulsions
III	Open fracture with extensive soft-tissue laceration (>10 cm), damage, or loss or an open segmental fracture. This type also includes open fractures caused by farm injuries, fractures requiring vascular repair, or fractures that have been open for 8 hours prior to treatment
IIIA	Type III fracture with adequate periosteal coverage of the fracture bone despite the extensive soft-tissue laceration or damage
IIIB	Type III fracture with extensive soft-tissue loss and periosteal stripping and bone damage. Usually associated with massive contamination. Will often need further soft-tissue coverage procedure (i.e. free or rotational flap)
IIIC	Type III fracture associated with an arterial injury requiring repair, irrespective of degree of soft-tissue injury

# 2022 Pediatric Open Fracture Evaluation Clinical Guideline

Fracture Classification	Initial Antibiotic/Dose
Gustilo Grade I or II	Cefazolin 50mg/kg (max 2000 mg)
Gustilo Grade III	Cefazolin 50mg/kg (max 2000 mg) <b>PLUS</b> Gentamicin 5mg/kg*
Open Extremity Fracture With Suspected Fecal/Clostridal Contamination (Farm-Related Injury)	Cefazolin 50mg/kg (max 2000 mg) <b>PLUS</b> Gentamicin 5mg/kg* <b>PLUS</b> Clindamycin 13mg/kg (max 600 mg)
Alternatives if Patient Has Severe Allergy to Above Antibiotics	<i>Beta-lactam Allergy:</i> Clindamycin 13mg/kg (max 600 mg) <i>Beta-lactam/Gentamicin Allergy:</i> Ertapenem 15mg/kg (max 1000 mg)

\*Consult ED Pharmacist for Appropriate Dosing Using Actual or Ideal Body Weights